

Mechanisms of Development 45 (1994) 285-286



Contents to Volume 45

No. 1		WJ. Park, J. Liu and P.N. Adler The frizzled gene of Drosophila encodes a membrane	
-	_	protein with an odd number of transmembrane domains	127
C. Hartmann, H. Taubert, H. Jäckle and M.J. Pankratz A two-step mode of stripe formation in the <i>Drosophila</i> blastoderm requires interactions among primary pair rule	1	R. Sugaya, S. Ishimaru, T. Hosoya, K. Saigo and Y. Emori A Drosophila homolog of human proto-oncogene ret tran- siently expressed in embryonic neuronal precursor cells including neuroblasts and CNS cells	139
genes	3	P.L. Pfeffer and E.M. De Robertis	
I.J. Mason, F. Fuller-Pace, R. Smith and C. Dickson FGF-7 (keratinocyte growth factor) expression during		Regional specificity of RAR γ isoforms in Xenopus development	147
mouse development suggests roles in myogenesis, fore- brain regionalisation and epithelial-mesenchymal interac- tions	15	M. Mével-Ninio, I. Guénal and B. Limbourg-Bouchon Production of dominant female sterility in <i>Drosophila melanogaster</i> by insertion of the ovo ^{D1} allele on autosomes: use of transformed strains to generate germline	
R.A. Cameron, L.C. Smith, R.J. Britten and E.H. Davidson Ligand-dependent stimulation of introduced mammalian		mosaics	155
brain receptors alters spicule symmetry and other morpho- genetic events in sea urchin embryos	31	K. Yoshida, I. Chambers, J. Nichols, A. Smith, M. Saito, K. Yasukawa, M. Shoyab, T. Taga and T. Kishimoto Maintenance of the pluripotential phenotype of embryonic	
E.M.L. Rabelo, B.S. Baker and J.R. Tata Interplay between thyroid hormone and estrogen in modulating expression of their receptor and vitellogenin genes		stem cells through direct activation of gp130 signalling pathways	163
during Xenopus metamorphosis	49	J.C. Richardson, A.M.G. Estrabot and H.R. Woodland	
T. Morio, I. Takeuchi and M. Tasaka Cooperation of positively and negatively acting promoter elements determines prespore-specific transcription of		XrelA, a Xenopus maternal and zygotic homologue of the p65 subunit of NF-κB. Characterisation of transcriptional properties in the developing embryo and identification of	172
Dp87 gene in Dictyostelium	59	a negative interference mutant	173
P. Cardellini, C. Polo and S. Coral			
Suramin and heparin: aspecific inhibitors of mesoderm induction in the <i>Xenopus laevis</i> embryo	73	No. 3	
No. 2			
Issue contents	89	Issue contents	191
P. Dollé, V. Fraulob, P. Kastner and P. Chambon Developmental expression of murine retinoid X receptor (RXR) genes	91	T. Thomas and M. Dziadek Expression of collagen $\alpha 1$ (IV), laminin and nidogen genes in the embryonic mouse lung: implications for branching morphogenesis	193
W. Lukowitz, C. Schröder, G. Glaser, M. Hülskamp and D. Tautz		C. Kaltschmidt, M. Muller, G. Brem and R. Renkawitz	
Regulatory and coding regions of the segmentation gene hunchback are functionally conserved between Drosophila virilis and Drosophila melanogaster	105	DNase I hypersensitive sites far upstream of the rat tryp- tophan oxygenase gene direct developmentally regulated transcription in livers of transgenic mice	203
J. Lef, J.H. Clement, R. Oschwald, M. Köster and W. Knöchel Spatial and temporal transcription patterns of the fork-		G.I. Gallicano, C.A. Larabell, R.W. McGaughey and D.G. Capco	
head related XFD-2/XFD-2' genes in Xenopus laevis embryos	117	Novel cytoskeletal elements in mammalian eggs are com- posed of a unique arrangement of intermediate filaments	211

morphogenesis

C. Mendelsohn, S. Larkin, M. Mark, M. LeMeur, J. Clifford, A. Zelent and P. Chambon RARβ isoforms: distinct transcriptional control by retinoic		R.G. Franks and S.T. Crews Transcriptional activation domains of the single-minded bHLH protein are required for CNS midline cell develop-	
acid and specific spatial patterns of promoter activity		ment	269
during mouse embryonic development	227		
		Author Index	279
D.R. Soprano, M. Gyda, III, H. Jiang, D.C. Harnish, K. Ugen,			
M. Satre, L. Chen, K.J. Soprano and D.M. Kochhar		Subject Index	281
A sustained elevation in retinoic acid receptor-\(\beta \) mRNA			
and protein occurs during retinoic acid-induced fetal dys-		Contents to Volume 45	285

243

255

P.S. Leahy, R.A. Cameron, M.A. Knox, R.J. Britten and E.H. Davidson Development of sibling inbred sea urchins: Normal em-

Development of sibling inbred sea urchins: Normal embryogenesis, but frequent postembryonic malformation, arrest and lethality

